



Standard 14: Triple Integrals - Cylindrical & Spherical Coord

Recall the last example from rectangular coordinates:

example. Evaluate $\iiint_E \sqrt{3x^2+3z^2} dV$ where E is the solid bounded by $y=2x^2+2z^2$ and the plane $y=8$.



$$D = \{(x,y) \mid x^2+y^2 \leq 4\}$$

$$\begin{aligned} \iiint_E \sqrt{3x^2+3z^2} dV &= \iint_D \left[\int_{2x^2+2z^2}^8 \sqrt{3x^2+3z^2} dy \right] dA \\ &= \int_{-4}^4 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} \int_{2x^2+2z^2}^8 \sqrt{3x^2+3z^2} dy dz dx \end{aligned}$$

The projection onto the xz -plane, the region D , is a disk. The equation of the disk comes from $8=2x^2+2z^2$. This region can be best described using something like polar coordinates for the xy -plane. Instead of the usual $x=r\cos\theta$ and $y=r\sin\theta$, we use $x=r\cos\theta$ and $z=r\sin\theta$. Thus

The region $D = \{(x,y) \mid x^2+y^2 \leq 4\}$ becomes $D = \{(r,\theta) \mid 0 \leq \theta \leq 2\pi, 0 \leq r \leq 2\}$. We also have to reflect these changes in the integral:

$$\begin{aligned} \iiint_E \sqrt{3x^2+3z^2} dV &= \int_{-4}^4 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} \int_{2x^2+2z^2}^8 \sqrt{3x^2+3z^2} dy dz dx \\ &= \int_0^{2\pi} \int_0^2 \int_{2r^2}^8 \sqrt{3r^2} \cdot r dz dr d\theta \end{aligned}$$

$\leftarrow dA = r dr d\theta$

cylindrical coordinates

This set of conversions is called cylindrical coordinates and is an extension of polar coordinates into three dimensions. In the example above we used the conversions for E 's in which the D is in the xz -plane, there is a different set for each type of E we have seen:

(i) $E = \{(x,y,z) \mid (x,y) \in D, u_1(x,y) \leq z \leq u_2(x,y)\}$ (ii) $E = \{(x,y,z) \mid (y,z) \in D, u_1(y,z) \leq x \leq u_2(y,z)\}$ (iii) $E = \{(x,y,z) \mid (x,z) \in D, u_1(x,z) \leq y \leq u_2(x,z)\}$

$$x = r\cos\theta$$

$$y = r\sin\theta$$

$$z = z$$

$$x^2+y^2 = r^2$$

$$dV = r dz dr d\theta$$

$$y = r\cos\theta$$

$$z = r\sin\theta$$

$$x = x$$

$$y^2+z^2 = r^2$$

$$dV = r dx dr d\theta$$

$$x = r\cos\theta$$


$$z = r\sin\theta$$

$$y = y$$

$$x^2+z^2 = r^2$$

$$dV = r dy dr d\theta$$

example. Set up $\iiint_E y dV$ where E is the region that lies below the plane $z=x+2$, above the xy -plane and between the cylinders $x^2+y^2=1$ and $x^2+y^2=4$.



$$x = r\cos\theta$$

$$y = r\sin\theta$$

$$z = z$$

$$0 \leq z \leq x+2$$

becomes

$$0 \leq z \leq r\cos\theta+2$$

$$D = \{(x,y) \mid 1 \leq x^2+y^2 \leq 4\}$$

$$= \{(r,\theta) \mid 1 \leq r \leq 2, 0 \leq \theta \leq 2\pi\}$$

$$\iiint_E y dV = \int_0^{2\pi} \int_1^2 \int_0^{r\cos\theta+2} r\sin\theta \cdot r dz dr d\theta$$

example. Convert the following integral into cylindrical coordinates: $\int_0^1 \int_0^{\sqrt{1-x^2}} \int_{-x}^{\sqrt{x^2+y^2}} xyz dz dy dx$.

Rectangular coordinate bounds:

$$0 \leq x \leq 1, 0 \leq y \leq \sqrt{1-x^2}, -x \leq z \leq \sqrt{x^2+y^2}$$

Cylindrical conversion:

$$x = r\cos\theta, y = r\sin\theta, z = z, dV = r dz dr d\theta$$

Sketch xy -plane:




Rewrite bounds:

$$0 \leq r \leq 1, 0 \leq \theta \leq \frac{\pi}{2}, -r\cos\theta \leq z \leq r$$

$$\int_0^{\pi/2} \int_0^1 \int_{-r\cos\theta}^r (r\cos\theta)(r\sin\theta)(z) \cdot r dz dr d\theta$$

spherical coordinates

There is another extension of polar coordinates into three dimensions; it is given by rotating polar coordinates. Spherical coordinates are (ρ, θ, φ) where ρ is the distance from the origin, θ is the angle made with the positive x -axis in the xy -plane, and φ is the angle made with the positive z -axis. Here is a visual of the conversion from rectangular to spherical:



$$x = \rho \sin(\varphi) \cos(\theta)$$

$$y = \rho \sin(\varphi) \sin(\theta)$$

$$z = \rho \cos(\varphi)$$

$$x^2 + y^2 + z^2 = \rho^2$$


$$dV = \rho^2 \sin(\varphi) d\rho d\theta d\varphi$$

with the restriction:

$$\rho \geq 0$$

$$0 \leq \varphi \leq \pi$$

example. Set up $\iiint_E z \, dV$ where E is inside both $x^2 + y^2 + z^2 = 2$ and the cone that makes an angle of $\pi/3$ with negative z -axis and has $x \leq 0$.



$$0 \leq \rho \leq \sqrt{2}$$

$$\frac{2}{3}\pi \leq \varphi \leq \pi$$

$$\frac{1}{2}\pi \leq \theta \leq \frac{3}{2}\pi$$

be careful when computing $\varphi \neq \theta$

they care about the positive axis

$$\iiint_E z \, dV = \int_{\frac{1}{2}\pi}^{\frac{3}{2}\pi} \int_{\frac{2}{3}\pi}^{\pi} \int_0^{\sqrt{2}} (\rho \cos(\varphi)) (\rho \sin(\varphi) \cos(\theta)) \rho^2 \sin(\varphi) d\rho d\theta d\varphi$$